




# Search for exotica at



- **Introduction**
- **A topical review of Exotica @ CMS**
- **Outlook and Conclusions**





Federal Ministry  
of Education  
and Research




Physics  
Institute III A

**RWTH**AACHEN  
UNIVERSITY

# Search for exotica at



- Introduction
- A topical review  
Exotica @ CMS
- Outlook and



Federal Ministry  
of Education  
and Research



Physics  
Institute III A

THAACHEN  
UNIVERSITY







# The Compact Muon Solenoid




Apr 26, 2013

LHC pp run in 2012:  
 $\sim 20 \text{ fb}^{-1}$  @  $\sqrt{s} = 8 \text{ TeV}$

Length:  $\sim 21 \text{ m}$   
Height:  $\sim 15 \text{ m}$   
Weight:  $12500 \text{ t}$

**CMS Integrated Luminosity, pp, 2012,  $\sqrt{s} = 8 \text{ TeV}$**

Data included from 2012-04-04 22:37 to 2012-12-16 20:49 UTC





# Searches for BSM (what are Exotica?)

- ▶ ... are a key part of the CMS physics program (and substantial)
- ▶ Supersymmetry, ...
- ▶ Technicolor, ...
- ▶ Extra dimensions, ...
- ▶ Extended gauge sector, 4<sup>th</sup> gen.
- ▶ Substructure, Leptoquarks
- ▶ Dark matter
- ▶ Black holes, unparticles
- ▶ (Non-)resonant structures
- ▶ Rare decays
- ▶ Metastable, long-lived particles
- ▶ Signature-based ↔ model-inspired
- ▶ Already more than 100 BSM physics papers!

**Theories**

**Concepts**

**Signatures**



# Searches for BSM (what are Exotica?)

- ▶ ... are a key part of the CMS physics program (and substantial)
- ▶ Supersymmetry, ...
- ▶ Technicolor, ...
- ▶ Extra dimensions, ...
- ▶ Extended gauge sector, 4<sup>th</sup> gen.
- ▶ Substructure, Leptoquarks
- ▶ Dark matter
- ▶ Black holes, unparticles
- ▶ (Non-)resonant structures
- ▶ Rare decays
- ▶ Metastable, long-lived particles
- ▶ Signature-based ↔ model-inspired
- ▶ Already more than 100 BSM physics papers!



© W.Sauber via wikimedia commons



# Searches for BSM (what are Exotica?)


- ▶ ... are a key part of the CMS physics program (and substantial)
- ▶ Supersymmetry, ...
- ▶ Technicolor, ...
- ▶ Extra dimensions, ...
- ▶ Extended gauge sector, 4<sup>th</sup> gen.
- ▶ Substructure, Leptoquarks
- ▶ Dark matter
- ▶ Black holes, unparticles
- ▶ (Non-)resonant structures
- ▶ Rare decays
- ▶ Metastable, long-lived particles
- ▶ Signature-based  $\leftrightarrow$  model-inspired
- ▶ Already more than 100 BSM physics papers!





# Searches for BSM (what are Exotica?)


- ▶ ... are a key part of the CMS physics program (and substantial)
- ▶ Supersymmetry, ...
- ▶ Technicolor, ...
- ▶ Extra dimensions, ...
- ▶ Extended gauge sector, 4<sup>th</sup> gen.
- ▶ Substructure, Leptoquarks
- ▶ Dark matter
- ▶ Black holes, unparticles
- ▶ (Non-)resonant structures
- ▶ Rare decays
- ▶ Metastable, long-lived particles
- ▶ Signature-based ↔ model-inspired
- ▶ Already more than 100 BSM physics papers!



# Summary of Exotica at CMS



## CMS EXOTICA 95% CL EXCLUSION LIMITS (TeV)





# Overview

- ▶ Monojets
  - ◆ Large extra dim., dark matter, ...
- ▶ Di-jet resonances (w, w/o b-tag)
  - ◆ Many interpretations
- ▶ 3-jet resonances (RPV SUSY)
- ▶ Di-lepton resonances
- ▶ Di-lepton large extra dimensions
- ▶ Single lepton + MET
  - ◆  $W'$ , Universal ED, CI, DM
- ▶  $W' \rightarrow tb$
- ▶ 2<sup>nd</sup> generation leptoquarks (l+jets)
- ▶  $Z' \rightarrow t\bar{t}$  searches
  - ◆ Semi-leptonic, all-hadronic
- ▶ Vector-like  $T' \rightarrow tZ, tH, bW$
- ▶ Vector-like  $B' \rightarrow tW, bH, bZ$
- ▶  $Q = 5/3$  top partners
- ▶ High mass di-boson resonances
  - ◆  $W'$ , extra dimensions, ...
- ▶ Microscopic black holes
- ▶ Jet extinction
  - ◆ Black holes, ...
- ▶ Displaced jets
  - ◆ Split/RPV SUSY, hidden valley, ...
- ▶ Heavy stable charged particles

# Overview

- ▶ Monojets
    - ◆ Large extra dim., dark matter, ...
  - ▶ Di-jet resonances (w, w/o b-tag)
    - ◆ Many interpretations
  - ▶ 3-jet resonances (RPV SUSY)
  - ▶ Di-lepton resonances
  - ▶ Di-lepton large extra dimensions
  - ▶ Single lepton + MET
    - ◆  $W'$ , Universal ED, CI, DM
  - ▶  $W' \rightarrow tb$
  - ▶ 2<sup>nd</sup> generation leptoquarks (l+jets)
  - ▶  $Z' \rightarrow t\bar{t}$  searches
    - ◆ Semi-leptonic, all-hadronic
  - ▶ Vector-like  $T' \rightarrow tZ, tH, bW$
  - ▶ Vector-like  $B' \rightarrow tW, bH, bZ$
  - ▶  $Q = 5/3$  top partners
  - ▶ High mass di-boson resonances
    - ◆  $W'$ , extra dimensions, ...
  - ▶ Microscopic black holes
  - ▶ Jet extinction
    - ◆ Black holes, ...
  - ▶ Displaced jets
    - ◆ Split/RPV SUSY, hidden valley, ...
  - ▶ Heavy stable charged particles
- : Next talk
- : Next-to-next talk







# Aside: $B_s \rightarrow \mu\mu$

BPH-13-007



- Highly suppressed in standard model (FCNC)
- Very sensitive to new physics in loops
- Finally observed by CMS and LHCb
- Result compatible with SM expectation







$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.9 \pm 0.7) \times 10^{-9}$$

# Aside: $B_s \rightarrow \mu\mu$

BPH-13-007



- ▶ Highly suppressed in standard model (FCNC)
- ▶ Very sensitive to new physics in loops
- ▶ Finally observed by CMS and LHCb
- ▶ Result compatible with SM expectation



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.9 \pm 0.7) \times 10^{-9}$$








# “Mono”-jets

EXO-12-048





- ▶ Versatile search channel for many “invisible” new particles
- ▶ Highly sensitive to extra dimensions, unparticles, dark matter, ...
- ▶ Main knobs: jet  $p_T$ , missing  $E_T$ 
  - ◆ Allow second jet (not back-to-back)









**Large extra dimensions:**  
**Fundamental Planck scale  $M_D > 3... 5 \text{ TeV}$**   
**(up to 5.7 TeV @ NLO)**




**(Stable, scalar) unparticles**

# “Mono”-jets

EXO-12-048





- Pair production of dark matter characterized by a contact interaction effective theory
- Limits translated to DM-nucleon cross sections in order to compare with direct detection experiments



- Many operators, validity range of effective theory, new models, ...



- Di-jet mass highly sensitive to many new physics models including excited quarks and contact interactions, axigluons,  $W'$ ,  $Z'$ , ...




- Present results in a way to allow straightforward application to new models


- Di-jet mass highly sensitive to many new physics models including excited quarks and contact interactions, axigluons,  $W'$ ,  $Z'$ , ...

Model	Final State	Obs. Mass Excl. [TeV]	Exp. Mass Excl. [TeV]
String Resonance (S)	qg	[1.20,5.08]	[1.20,5.00]
Excited Quark ( $q^*$ )	qg	[1.20,3.50]	[1.20,3.75]
$E_6$ Diquark (D)	qq	[1.20,4.75]	[1.20,4.50]
Axigluon (A)/Coloron (C)	$q\bar{q}$	[1.20,3.60] + [3.90,4.08]	[1.20,3.87]
Color Octet Scalar ( $s_8$ )	gg	[1.20,2.79]	[1.20,2.74]
$W'$ Boson ( $W'$ )	$q\bar{q}$	[1.20,2.29]	[1.20,2.28]
$Z'$ Boson ( $Z'$ )	$q\bar{q}$	[1.20,1.68]	[1.20,1.87]
RS Graviton (G)	$q\bar{q}+gg$	[1.20,1.58]	[1.20,1.43]

CMS  
CMS Experiment at LHC, CERN  
Data recorded: Fri Oct 5 12:29:33 2012 CEST  
Run/Event: 204541 / 52508234  
Lumi section: 32



CMS  
CMS Experiment at LHC, CERN  
Data recorded: Fri Oct 5 12:29:33 2012 CEST  
Run/Event: 204541 / 52508234  
Lumi section: 32




Di-jet mass = 5.15 TeV

- Present results in a way to allow straightforward application to new models



# Di-jet searches ( $b\bar{b}$ , $bg$ )



EXO-12-023



Extend to  
(0,) 1, 2  
b-tags

- ▶ Benchmark: pair-produced gluinos decaying to 3 jets in RPV SUSY
- ▶ Combine 6 highest  $p_T$  jets into 20 unique triplet combinations
- ▶ To suppress wrong combinations and QCD, only accept triplets that satisfy
- ▶ Look for bump in falling spectrum

$$M_{jjj} < \sum_{i=1}^3 |p_{T|i} - \Delta$$







# Pair-produced 3-jet resonances


EXO-12-049



- ▶ In addition to inclusive search, apply b-tagging for enhanced sensitivity to decays with b quarks (gluino  $\rightarrow$  udb, csb)
- ▶ Inclusive search: exclude RPV decaying gluinos with  $M < 650$  GeV
- ▶ b-tagged: exclude  $200 < M < 835$  GeV



- New heavy gauge bosons  $\rightarrow$  narrow  $ee, \mu\mu$  resonances in the TeV region



$\rightarrow$  Upper limit on the ratio of  $Z'$  (or  $G_{KK}$  or ...) to SM  $Z$  production

95% CL mass limits:



$Z'$  SSM

$Z'$  Psi

2.96 TeV

2.60 TeV

- New heavy gauge bosons  $\rightarrow$  narrow  $ee, \mu\mu$  resonances in the TeV region



95% CL mass limits:

$Z'_{\text{SSM}}$

2.96 TeV





$Z'_{\Psi}$

2.60 TeV



# Large extra dimensions (ADD) in $ee, \mu\mu$


- Virtual graviton production modifies Drell-Yan spectrum
- Signal region:  $M_{\mu\mu} > 1.8 \text{ TeV}$



$M_s$ (ADD) at LO 95% CL limits	Lumi. [fb <sup>-1</sup> ]	$\delta=3$ Exp.	$\delta=3$ Obs.	$\delta=6$ Exp.	$\delta=6$ Obs.	$\Lambda_T$ (GRW) [TeV]
CMS dimuon	20.6	4.34	4.33	3.07	3.06	3.64
CMS dielectron	19.6	4.62	4.64	3.27	3.28	3.90
Combined:	20.6+19.6	4.76	4.77	3.37	3.37	4.01

# Large extra dimensions (ADD) in $ee, \mu\mu$

EXO-12-027/031



- ▶ UV cut-off parameter  $M_s$ , not directly comparable to fundamental Planck scale  $M_D$
- ▶ Complementary to mono-jets and mono-photons
- ▶ Large part of interesting parameter space excluded

$M_s$ (ADD) at LO 95% CL limits	Lumi. [fb <sup>-1</sup> ]	$\delta=3$ Exp.	$\delta=3$ Obs.	$\delta=6$ Exp.	$\delta=6$ Obs.	$\Lambda_T$ (GRW) [TeV]
CMS dimuon	20.6	4.34	4.33	3.07	3.06	3.64
CMS dielectron	19.6	4.62	4.64	3.27	3.28	3.90
Combined:	20.6+19.6	4.76	4.77	3.37	3.37	4.01

# Single lepton + MET ( $e\nu, \mu\nu$ )

EXO-12-060






## ► Many models, e.g.:

- ◆ sequential SM with and w/o interference, no decays to WZ
- ◆ Universal extra dimensions
- ◆ Four fermion contact interactions

## ► In events with isolated $e$ or $\mu$ plus MET, use binned likelihood in $M_T$



$$M_T = \sqrt{2 \cdot p_T^\ell \cdot E_T^{\text{miss}} \cdot (1 - \cos \Delta\phi_{\ell, \nu})}$$





# Single lepton + MET ( $e\nu, \mu\nu$ )

EXO-12-060



# Single lepton + MET ( $e\nu, \mu\nu$ )

EXO-12-060



## ► Limits in SSM:



- ◆  $M > 3.35$  TeV (no interference)
- ◆  $M > 3.10$  TeV /  $3.60$  TeV (destr. / constr.)

## ► At high masses, $W'$ mostly off-peak

## ► Limits on HNC CI:


- ◆  $\Lambda > 13.0$  TeV (e) /  $10.9$  TeV ( $\mu$ )

Model	Channel	Observed limit	Expected limit
SSM	e	$m_{W'} < 3.20\text{TeV}$	$m_{W'} < 3.25\text{TeV}$
SSM	$\mu$	$m_{W'} < 3.15\text{TeV}$	$m_{W'} < 3.10\text{TeV}$
SSM	combined	$m_{W'} < 3.35\text{TeV}$	$m_{W'} < 3.40\text{TeV}$
SSMO	e	$m_{W'} < 3.60\text{TeV}$	$m_{W'} < 3.60\text{TeV}$
SSMO	$\mu$	$m_{W'} < 3.05\text{TeV}$	$m_{W'} < 3.30\text{TeV}$
SSMO	combined	$m_{W'} < 3.60\text{TeV}$	$m_{W'} < 3.60\text{TeV}$
SSMS	e	$m_{W'} < 3.00\text{TeV}$	$m_{W'} < 3.10\text{TeV}$
SSMS	$\mu$	$m_{W'} < 2.80\text{TeV}$	$m_{W'} < 2.90\text{TeV}$
SSMS	combined	$m_{W'} < 3.10\text{TeV}$	$m_{W'} < 3.20\text{TeV}$
$W_{KK}^2$	$\mu=0.05$ TeV, combined	$m_{W_{KK}^2} < 1.7\text{TeV}$	$m_{W_{KK}^2} < 1.7\text{TeV}$
$W_{KK}^2$	$\mu=10.0$ TeV, combined	$m_{W_{KK}^2} < 3.7\text{TeV}$	$m_{W_{KK}^2} < 3.6\text{TeV}$
HNC CI	e	$\Lambda < 13.0$ TeV	$\Lambda < 13.3$ TeV
HNC CI	$\mu$	$\Lambda < 10.9$ TeV	$\Lambda < 12.2$ TeV







# Single lepton + MET ( $e\nu, \mu\nu$ )

EXO-13-004






- ▶ Dark matter production:  $W$  recoiling against pair-produced dark matter
- ▶ Reinterpretation of leptonic  $W'$  search
- ▶ Consider vector and axial-vector like couplings
- ▶ Possible interference effects
- ▶ First limits on “monolepton” DM








- Searches in  $W' \rightarrow$  quark final states important to complement leptonic searches, in case decays into leptons are suppressed
  - Also, can reconstruct mass (modulo ambiguities)
- Enhanced couplings to 3<sup>rd</sup> generation possible (and easier!)
- Signature is a high  $p_T$  isolated lepton, large MET, and 2 b-jets (one b-tag required)
- Exclude  $M(W') < 2.03 \text{ TeV}$  at 95% CL for  $W_R$  and  $W_L$  w/o interference
- Limits for arbitrary combinations of LH or RH couplings





- ▶ Carrying both lepton and baryon numbers, coupling to lepton-quark pair
- ▶ Fractional charge, three generations
- ▶ Here: pair production of scalar LQ, coupling to one generation only
- ▶ Parameters: LQ mass, BF to  $lq$  ( $\beta$ )
- ▶ Optimize cuts for each hypothetical  $M(\text{LQ})$ 
  - ◆  $S_T$  ;  $M(\mu\mu)$ ,  $M_T(\mu\nu)$  ;  $M(\mu, \text{jet})$
  - ◆  $S_T$  = scalar  $p_T$  sum of  $\mu$ ,  $\mu$  ( $\nu$ ),  $j1$ ,  $j2$



# LQ2 ( $\mu\nu+jj$ , $\mu\mu+jj$ )

EXO-12-042



**$M > 1070 \text{ (785) GeV}$   
for  $B(LQ \rightarrow \mu q) = 1 \text{ (0.5)}$**



# $Z' \rightarrow t\bar{t} \rightarrow e/\mu + \text{jets}$

B2G-12-006

B2G-13-001







► 1 lepton + MET +  $\geq 2$  jets

► 0 or 1 b-tag

► Define  $\chi^2$  for top hypotheses

- ◆ Separately optimized for low mass ( $M_{t\bar{t}} < \sim 1$  TeV, non-boosted) and high mass (boosted) regime



# $Z' \rightarrow t\bar{t} \rightarrow e/\mu + \text{jets}$

B2G-12-006

B2G-13-001






## ► Topcolor $Z'$ limits with 95% CL:

◆ Narrow:  $M(Z') > 2.1 \text{ TeV}$

◆ Wide:  $M(Z') > 2.7 \text{ TeV}$

## ► Randall-Sundrum model:

◆  $M(\text{KK gluon}) > 2.5 \text{ TeV}$



# $Z' \rightarrow t\bar{t}$ all hadronic



B2G-12-005

B2G-13-001



- ▶ Boosted top- and W-tagging
- ▶ Topcolor  $Z'$  limits with 95% CL:
  - ♦ Narrow:  $M(Z') > 1.7 \text{ TeV}$
  - ♦ Wide:  $M(Z') > 2.35 \text{ TeV}$
- ▶ Randall-Sundrum model:
  - ♦  $M(\text{KK gluon}) > 1.8 \text{ TeV}$

## Combination





# Vector-like $T' \rightarrow tZ/tH/bW$

B2G-12-015



- ▶ Combined information from single lepton, SS and OS di-lepton, tri-lepton
- ▶ Bin by W-tags, N(jets), N(b-jets),  $H_T$ , MET, lepton  $p_T$ , 3<sup>rd</sup> / 4<sup>th</sup> jet  $p_T$
- ◆ OS targeting  $tZtZ$ : on-Z,  $\geq 5$  jets,  $\geq 2$  b-jets,  $H_T > 500$  GeV,  $S_T > 1000$  GeV
- ◆ OS targeting  $bWbW$ : off-Z, 2-3 jets,  $H_T > 300$  GeV,  $S_T > 900$  GeV
- ◆ SS targeting  $tZ$  or  $tH$ :  $\geq 3$  jets,  $H_T > 500$  GeV,  $S_T > 700$  GeV, lepton flavor cat.eg.s.
- ◆ Tri-lepton targeting  $tZ$  or  $tH$ :  $\geq 3$  jets,  $H_T > 500$  GeV,  $S_T > 700$  GeV, lepton flavor categories






# Vector-like $T' \rightarrow tZ/tH/bW$

B2G-12-015




- ▶ Combine all channels to get limits
- ▶ As function of branching fractions, exclude masses between 687 and 782 GeV

CMS preliminary  $\sqrt{s} = 8 \text{ TeV}$   $19.6 \text{ fb}^{-1}$



CMS preliminary  $\sqrt{s} = 8 \text{ TeV}$   $19.6 \text{ fb}^{-1}$






# Vector-like $b' \rightarrow tW/bH/bZ$ (lepton+jets)

B2G-12-019



- ▶ One electron/muon, MET,  $\geq 4$  jets including 1 with b-tag
- ▶ Classify events based on “V-tags” = number of jets consistent with boosted W, Z, or H boson
- ▶ Search for excess in  $S_T$  = scalar sum of jet  $p_T$ , lepton  $p_T$ , MET
- ◆ Fit to background and signal templates in different categories





# Vector-like $b' \rightarrow tW/bH/bZ$ (lepton+jets)

B2G-12-019



- ▶ As a function of  $b' \rightarrow tW, bH, bZ$  branching fractions, exclude up to  $M(b') < 732 \text{ GeV}$
- ▶ Highest sensitivity for  $b' \rightarrow tW$






# Vector-like $b' \rightarrow bZ$ (di-lepton + jets)

B2G-12-021




- Consider decay to  $bZ$  and  $tW$
- Reconstruct resonance of  $e^+e^-$  and  $\mu^+\mu^-$  pairs compatible with  $Z$ , plus  $b$ -jet
- Assuming  $B(b' \rightarrow bZ) = 100\%$ , exclude  $M(b') < 700 \text{ GeV}$
- ... plus limits on  $B(b' \rightarrow bZ)$  vs.  $M(b')$









$$T_{5/3} \rightarrow t W$$

B2G-12-012



- ▶ Composite Higgs or models with extra dimensions
- ▶ Same-sign di-leptons (e or  $\mu$ )
- ▶ Allow for boosted top and W
- ▶ 5 or more constituents (top=3, W=2) in addition to the two leptons
- ▶  $H_T > 900$  GeV  $\rightarrow$  data: 11 events; bgrd:  $6.6 \pm 2.0$
- ▶ Exclude  $M(T_{5/3}) < 770$  GeV at 95% CL




# High mass di-boson resonance searches




## EXO-12-025:

$$W'/\rho_{TC} \longrightarrow WZ \longrightarrow 3l + MET$$




## EXO-12-021:

$$G_{bulk} \longrightarrow WW \longrightarrow l + jet + MET$$




## EXO-12-022:

$$G_{bulk} \longrightarrow ZZ \longrightarrow 2l + 2jets$$



## EXO-12-024:

$$G_{RS} \longrightarrow WW/ZZ, W' \longrightarrow WZ$$






# $W' / \rho_{TC} \rightarrow WZ \rightarrow 3l + MET$

EXO-12-025



- Use  $M_{WZ}$  (taking MET into account) and  $\Sigma p_T(l)$
- Modified ID and isolation for muons (from boosted Z)
- $W' \rightarrow WZ$  excluded for  $0.17 < M < 1.45$  TeV







## ► $G_{RS} \rightarrow WW, WZ$ and $W' \rightarrow WZ$ in di-jets, with $W$ and $Z \rightarrow jj$

- ♦ Jets from  $W/Z$  decays boosted and merged into single jets
- ♦ Each jet required to pass “W/Z-tagger”: pruned jet mass, N-subjettiness

## ► Exclude

- ♦  $G_{RS} (k/M_{PL}=0.1) \rightarrow WW(ZZ)$  for  $1.0 < M < 1.59$  (1.17) TeV
- ♦  $W' \rightarrow WZ$  for  $1.0 < M < 1.73$  TeV
- ♦  $q^* \rightarrow qW (qZ)$  for  $1.0 < M < 3.23$  (3.00) TeV





# Black holes



Google

helden - wenn dein land dich braucht rtl



Web

**Bilder**

Maps

Shopping

Mehr ▾

Suchoptionen

Sat





“Helden” (Heroes) last week on German television (RTL)

# Microscopic black holes


EXO-12-009



- ▶ Would decay to multiple objects: jets, leptons, photons, ...
- ▶ Distribution of interest: scalar  $p_T$  sum =  $S_T$
- ▶ For backgrounds, extrapolate shape from low ( $N=2$ ,  $1.8 < S_T < 2.8$  TeV) to high multiplicity
- ◆ Normalization from low  $S_T$ ,  $1.9 < S_T < 2.3$  TeV



.....




# Microscopic black holes



EXO-12-009





- Derive model-dependent limits, as well as limits on excess  $S_T$  for different object multiplicities



13 jets,  $S_T = 4.5$  TeV



- ▶ Even if black holes (or similar phenomena) are produced, we may miss them (trigger, no decay to our “objects”, not yet significant, ...)
- ▶ However, above production threshold, SM processes are highly suppressed
  - ◆ Good place to check are inclusive jets
  - ◆ PDF and jet energy scale uncertainty are important




- ▶ Limit on the extinction energy scale (fundamental Planck scale)  **$M > 3.3 \text{ TeV}$**







- ▶ Massive long-lived particles decaying to displaced jets can occur in many models: split SUSY, RPV SUSY, hidden valley models etc.
- ▶ Benchmark here is a heavy scalar:  $gg \rightarrow H^0 \rightarrow X^0 X^0 \rightarrow qq\, qq$ 
  - ◆ Mean decay length of  $X^0$ : 3... 300cm
- ▶ Search for di-jets from a common, displaced vertex
- ▶ Background suppression based on vertex track multiplicity, fraction of tracks with positive  $d_0$ , likelihood discriminant



2 candidate events,  
compatible with  
background  
expectation

- ▶ Massive long-lived particles decaying to displaced jets can occur in many models: split SUSY, RPV SUSY, hidden valley models etc.
- ▶ Benchmark here is a heavy scalar:  $gg \rightarrow H^0 \rightarrow X^0 X^0 \rightarrow qq\, qq$ 
  - ◆ Mean decay length of  $X^0$ : 3... 300cm
- ▶ Search for di-jets from a common, displaced vertex
- ▶ Background suppression based on vertex track multiplicity, fraction of tracks with positive  $d_0$ , likelihood discriminant



Limits in the range  
0.3 ... 300 fb

# Heavy stable charged particles

EXO-12-026



- Comprehensive analysis using time-of-flight (muon system) and/or dE/dx (tracker)





◆ Lifetime  $> 1$  ns, mass  $> 100$  GeV

- Make use of

◆ Track  $p_T$ : inner tracker

◆ Muon  $1/\beta$ : muon system




◆ Track  $I_{as}$ : dE/dx MIP incompatibility





# Heavy stable charged particles

- Results for long-lived gluinos, scalar top quarks, scalar tau leptons, fractionally/multiply charged leptons
- Mass limits up to  $> 1300$  GeV (gluinos)







# Outlook

- ▶ Run 2 from 2015 to 2017:  $\sqrt{s} = 13 - 14$  TeV,  $\int L dt \sim 40 - 45 \text{ fb}^{-1}/\text{a}$
- ▶ Longterm HL-LHC from  $\sim 2023$ ,  $\int L dt \sim 300 \text{ fb}^{-1}/\text{a}$ ,  **$3000 \text{ fb}^{-1}$  total**,  $\sim 140$  PU
- ▶ Example: vector-like  $T' \rightarrow tZ, tH, bW$  (as B2G-12-015)

FTR-13-026



5 $\sigma$  discovery reach up to  $M(T')$  of 1500 GeV  
(compared to  $\sim 700$  GeV exclusion @8 TeV)


- ▶ ECFA workshop last week:  
<https://indico.cern.ch/conferenceDisplay.py?confId=252045>

# Conclusions



- ▶ No stone left unturned in BSM searches at the LHC and CMS
- ▶ While there's an overwhelming output of results, many signatures are not yet fully exploited
  - ◆ New physics may still be hiding in the data already collected
  - ◆ Important guidance from the “new boson”
- ▶ Unfortunately no compelling evidence for new physics
  - ◆ But many results exploring new territory
- ▶ Many publications with full 2012 data set close to publication
- ▶ Preparing for next run in 2015
- ▶ This and more at

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsEXO> and [PhysicsResultsB2G](https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsB2G)



<http://www.hep.physik.uni-siegen.de/~gruppen>

**Backup**

# Summary

